

# Using fennel and its oil as a preservative and functional food to produce food and Drink products to be used to alleviate cough symptoms

Jehan I. Saber<sup>1</sup>, Dalia H. Eshra<sup>2</sup>

## ABSTRACT

Fennel seeds and its aromatic oil contain ingredients which are active biologically with high nutritional value, as well as their antimicrobial and fungal effects and alleviation of some diseases symptoms. The aim of this research was to determine the physical properties of fennel seeds and its aromatic oil and their effect on the growth of microbes and alleviation of respiratory diseases symptoms such as cough and sore throat.

Preparation of fennel seeds drink and its aromatic oil at different ratios were used to treatment a group of individuals have cough and compare them with the control group. In addition, biscuits containing different ratios of fennel seeds and its aromatic oil were prepared.

The results of this study showed that fennel seeds contained a high percentage of protein, dietary fiber, carbohydrates, minerals, vitamins and flavonoids. The results also showed that fennel seeds and its aromatic oils contained substances, which have an effect in relieving cough and sore throat. Extract of methanol from fennel seeds has antioxidant and antimicrobial activities. This study also showed that biscuits containing fennel seeds and its aromatic oil can be stored for longer periods than biscuits without fennel seeds and its aromatic oil no change in sensory properties or growth of microbial during storage. Therefore, we can use it as a natural preservative. In addition, biscuits prepared containing the different percentages of fennel seed and its aromatic oil and were accepted by the panelists when they were stored. Its sensory properties were well maintained. The results also showed that drink of fennel and its oil had an effective effect in the treatment of cough.

**Keywords:** *Fennel, foods products, antibacterial activity, fennel oil aromatic.*

## INTRODUCTION

Nutraceutical is the harmonious sequence of 2 words “nutrition and pharmaceutical” that is defined as the food product that helps to reinforce health status along with medicinal benefits, comprehensive the treatment of diseases with safeguard. Many types of products that link under the group of nutraceuticals are dietary complement and functional foods (Chaturvedi *et al.*, 2011 & Bukhari *et al.*, 2014). According to IFIC (International Food Information Council), functional foods are defined as dietary components in which health avails are along with the

required nutrition. Many foods like breads, cereals, herbalists, medicinal and aromatic herbs, snack foods, prepared meals and many others are included in functional foods (Keservani *et al.*, 2010).

Food spoilage caused by microorganisms are of the most important trouble facing the food industry and consumers, so synthetic additives was used to inhibit growth of microorganisms (Al-Reza *et al.*, 2010 & Bajpai *et al.*, 2012). Because of consumer's fears

of the safety of foods containing synthetic chemicals, there is an increase interest in using natural antibacterial products for food preservation (Bajpai *et al.*, 2012). On the other hand, the long-term use of drugs to treat diseases has various adverse effects, so there is a need to use natural substances that have little side effects instead of using chemical treatments (Choi and Hwang, 2004).

Recently, awareness increased concerning spice beneficial physiological functions and its antimicrobial activity. In addition, use of essential oils as antimicrobial agents in food systems considered as determinant to increase the safety and increase shelf life of foods (Salgueiro *et al.*, 2010). And they are commonly used as a natural remedy for some diseases (Choi and Hwang, 2004).

Fennel (*Foeniculum vulgare*) is a familiar herb, it is universally known as Fennel and is known by more than 100 names and recognized as aromatic, herbaceous, and is widely cultivated and used as a culinary spice (Gori *et al.*, 2012 & Diao *et al.*, 2014). Fennel and its essential oil are used as flavoring in food products such as bread, pastries, and cheese consumed daily, in the raw form as salads and, boiled, grilled, in several dishes and even used in the preparation of herbal teas. They are also used as a constituent in pharmaceutical products to cure certain diseases and the inhibition of pain (Barros *et al.*, 2010 & He and Huang, 2011 & Rather *et al.*, 2012). It is traditional and popular herb and has a long history of use as a medicine. Studies showed that Fennel and its essential oil effectively control numerous infectious disorders of bacterial, fungal, viral, antimicrobial, and used to treat tuberculosis and respiratory diseases (Corona *et al.*, 2008 & Badgujar *et al.*, 2014). The antioxidant and antimicrobial characteristics of the fennel and its essential oil were exploited in products as

DOI: 10.21608/ASEJAIQJSAE.2019.44629

<sup>1</sup>Home Economics Dept., Fac. of Agric, Alex University El-Shatby, 21545, Alex., Egypt .Food

<sup>2</sup>Science & Technology Dept., Fac. of Agric., Alex. University, El-Shatby, 21545, Alex, Egypt.

Received June 13, 2019, Accepted August 07, 2019

natural preservatives (Telci *et al.*, 2009 & Diao *et al.*, 2014).

Thus, this work was undertaken to study the chemical component of fennel seeds, its oil, biological compounds, and their effect on inhibiting the growth of microorganisms in some food products and the use of fennel seeds and its oil in the treatment and relieve symptoms of respiratory diseases such as cough, laryngitis.

## MATERIALS AND METHODS

Fennel (*Foeniculum vulgare*) and its essential oil were procured from the local market of Alexandria, Egypt. Bacterial and fungal strains including: *Staphylococcus aerus* 29123, *Escherichia coli* 3518, *C. albicans Spp* and *Aspergillus niger* CAIM 147 were obtained from Dairy science Dep. Alex. University.

### Technological processes

The traditional methods were applied to prepare three samples of biscuits were prepared by different percentages of Fennel seeds and its essential oil as follows:

25% of Fennel seeds was added to the first sample, 50 % of Fennel seeds was added to the second sample, and 25 % Fennel seeds + 2 % of its essential oil (equivalent 2 -3 pt of oil) was added to third sample. Percentage of the seeds and its oil were calculated from the flour weigh. Moreover, the products were stored at room temperature for three months.

### Drinks of Fennel seeds and its oil

Teaspoon of Fennel seeds (equal 5 g) was added to small cup (150 ml) of water and boiled for 3 minutes (which is the perfect time to obtain the highest amount of antioxidants). The drink was left until it became warm, one drop of oil added to the mixture (Malhotra, 2012 & Kontogiorgis *et al.*, 2016). The same procedure was repeated using 10 g and 15 g of Fennel seeds. Persons suffering of cough had repeated this drink 4 to 5 times daily until symptoms disappear (7 to 3 days).

### Analytical methods

#### Proximate chemical composition

Moisture, crude protein, crude ether extract and total ash of Fennel seeds were determined according to AOAC (2006) unless otherwise stated. Carbohydrate content were calculated by difference.

#### Mineral determination

Mineral content of Fennel seeds were determined by atomic absorption spectrometry, flame photometry and spectrophotometry according to the methods of AOAC (2006).

### Gas Chromatography–Mass Spectrometry (GC–MS)

Oil of fennel seeds was analyzed on a gas chromatograph, coupled with mass spectrometer. Identification of different constituents of fennel oil was determined by the spectrum fragmentation pattern compared to the chromatograms from standards. Furthermore, the four major components of the fennel essential oils, trans-anethole, fenchone, estragole and limonene, were quantified by mean of the internal standard addition method (Mansour *et al.*, 2010 & Burkhardt *et al.*, 2015).

### Preparation of Fennel Seed Extracts

Fennel seeds powder were soaked in 70% ethanol (1:40, w/v) and left for 72 h at room temperature, shaking the mixture every now and then. The extract was filtered and the precipitate was re-extracted by the same process using the same solvent until the extraction was exhausted. The combined extracts were separately filtered through a filter paper. The filtered was dried under reduced pressure at 50 °C using a rotary vacuum evaporator. The crude extract was weighed and kept in a tightly closed container protected from light (Anwar *et al.* 2009).

### Antioxidant capacity by DPPH assay

Free radical scavenging activity were measured by using the 2,2- diphenyl-1-picrylhydrazil (DPPH) according to a modified method by Roby *et al.*, (2012). The effect of antioxidant on DPPH radical scavenging can be due to their hydrogen donating ability or radical scavenging activity. When a solution of DPPH is mixed with a substance that can donate a hydrogen atom, it then leads to a loss of this violet color.

The effect of Fennel seeds extract on (DPPH) radical was estimated using a solution of DPPH (0.135 mM) was prepared and 1 ml of this solution was mixed with 1 ml of the extract. The reaction mixture was left in the dark at room temperature for 30 min. The absorbance of the mixture was measured at 517 nm using butylated hydroxyl anisole (BHA) as a control .The radical scavenging activity is expressed by the following formula:

$$\text{DPPH inhibition \%} = \frac{\text{Initial absorbance} - \text{sample absorbance}}{\text{Initial absorbance}} \times 100$$

The half maximal inhibitory concentration (IC<sub>50</sub>) values denoted the concentration of sample required to scavenge 50% of DPPH free radicals.

### Total phenolic content

Total phenolic content was determined using folin Ciocalteu reagent. 1 ml of extract was mixed with 5 ml folin – Ciocalteu reagent and 4 ml of sodium carbonate (75 g/l). The mixture for 15 s and allowed to stand for

30 min for colour development. The absorbance was read at wavelength 765-nm.using spectrophotometer. Total phenolic content was expressed as mg gallic acid equivalent /g of extract (Ahmad *et al.*,2018).

#### Total flavonoid content

The total flavonoid content was determined using the aluminum chloride assay through colorimetry (Ahmad *et al.* ,2018). The absorbance of the reaction was measured at 500 nm with a UV- Visible spectrophotometer after 15 min. Distilled water was used as blank. Quantification was done based on a standard curve of rutin and expressed in mg of rutin equivalents / g of extract.

#### Antimicrobial, antibacterial and antifungal activities

Detected antimicrobial activity of Fennel seed extracts and its essential oil by selected two species of bacteria, one of them was Gram-negative bacteria (*Escherichia coli* 3518) and the other Gram-positive bacteria (*Staphylococcus aureus* 29123). in addition to one species of yeast (*Candida albicans* Spp) and one species of mold (*Aspergillus niger* CAIM 147), determined using the disk expansion assay. Isolates were cultured at 37°C for 24h in Nutrient Agar and bacterial suspensions were all set with Nutrient Broth to match McFarland standard No. 0.5 turbidity (Kaskatepe *et al.*, 2016). The antimicrobial activity was estimated by agar well diffusion method, a well was prepared in the plates with the help of a cork-borer (0.90 cm). The disc diameter was 6 mm (Whatman No. 1) paper. Different dilutions of the extracts and essential oils were made with methanol. Under aseptic conditions, the discs were placed on the agar plates and then take 7.5, 10, 12.5, 15 and 20 <sup>1</sup>-g from each of the extracts and essential oil dilutions were put on the discs. The plates were then incubated at 37 °C for bacteria and 25 °C for yeast and fungi for 24–48 h to get microbial growth. Diameters of microbial inhibition zones (mm) were measured and recorded (Roby *et al.* 2012).

#### Sensory evaluation

Colour, taste, odour, texture, (consistency) and overall acceptability of the products prepared containing the different percentages of Fennel extracts and its essential oil were assessed using 15 of arbitrators. The panelists were asked to score the above attributes according to a standard hedonic rating score from 9 (like extremely) to (dislike extremely): 1 = dislike extremely, 2 = dislike very much, 3 = dislike moderately, 4 = dislike slightly, 5= like/dislike, 6 = like slightly, 7 = like moderately, 8 = like very much, 9 = like extremely. as described by (Wichchukita and O'Mahony, 2014).

#### Sample of individuals

Twenty members of individuals were selected with cough and sore throat (10 smokers and 10 non-smokers), their age ranged from 25 to 40 years they were divided into two groups each group contains equal numbers of smoker and non-smokers:

- he first group (control): Ten people taking medicines to treat cough and sore throat (they was took treatment of cough which was recommended by doctor).
- cond group: (test group): Ten people who drink fennel seeds and its drink oil.

#### Statistical analysis:

Data were analysed using SPSS statistical analysis software package (Version 21).

### RESULTS AND DISCUSSION

#### Chemical composition and dietary fiber content of Fennel

Consumers have been more conscious of food related health troubles and using healthy food to Provide the body with important natural nutrients and resistance to some diseases. Table (1) shows the proximate chemical composition. These analyses have an important role in chemical screening of compounds. Proximate composition of fennel seed contains moisture, crude protein, crude fat, crude fiber, ash as (8.04, 10.18, 10.71, 18.01 and 12.87) % respectively and total carbohydrate by difference was 40.19% .

**Table 1. Proximate Composition of the Fennel seeds**

Component	% Value**
Moisture	8.04±0.23
Crude protein	10.18±0.39
Crude Fat	10.71±0.32
Crude fiber	18.01±0.73
Total ash	12.87±0.41
Carbohydrate*	40.19 ± 1.24

\* calculated by difference

\*\* Mean value ± S.D. on dry weight basis

**Table 2. Mineral content of Fennel seeds**

Mineral*	Amounting (mg/100g)
K	849.45
Na	15.91
Fe	9.69
Ca	583.6
Zn	2.89
Mg	85.87
Mn	209.35
P	470
Pb	ND

\* On dry weight basis

ND: Non detected

**Table 3. Relative percentage composition of fennel essential oil**

Compound	Relative Contents(%)	Compound	Relative Contents(%)
$\beta$ -cymene	0.30	Cis-anethole	0.29
D-limonene	4.44	Trans-anethole	77.91
Eucalyptol	0.23	Anisaldehyde	4.54
$\gamma$ -terpinen	0.84	Apiole	0.39
Anisole	3.59	Estragol	5.65
Camphor	0.46	<i>p</i> -acetonylanisole	0.59

This study are in harmony with investigations of (Badgujar *et al.*, 2014 & Bukhari *et al.* 2014) found that percentage of moisture, protein, fat, fiber and ash in fennel was 7.27, 9.5, 10.18, 18.97 and 13.4%, respectively. However, in this study was the moisture, protein, fat, and fiber content was lower than the values reported by (Malhotra 2012 & USDA. 2018). While, the ash content in this study is relatively higher than the values reported by (USDA. 2018).

#### Mineral contents of Fennel seeds

Mineral content of Fennel seeds were determined and presented in (Table 2).

Table (2 ) shows that fennel Seed contained iron (6.33mg/g), Zinc (2.89 mg/100g), Calcium (583.6 mg/100g), Manganese (209.35 mg/100g), Magnesium (83.87mg/100 g), sodium (15.91mg/100 g), Potassium (849.45 mg/100g) , phosphorus (470 mg/100g). These results agree with these reported by (Badgujar *et al.*, 2014 & Bukhari *et al.* 2014) The data obtained in the present study are more or less in accordance with those reported by (Koudela and Petříková 2008). Fennel seeds may be considered as a good source of K, Ca and Mg.

#### Chemical composition of the essential oil

The chemical composition of fennel oil is shown in Table 3. Fennel oil contains Trans-anethole (77.91%) this component was the highest, this is agreement with reported by (Mansour *et al.*, 2011 & Qiu *et al.*, 2012 & Ahmad *et al.*, 2018) the percentage of Trans-anethole was (65.28%, 88.91% and 70.72 % respectively. while, The other components were low.

**Table 4. Antioxidant activities using DPPH**

Parameter	Value <sup>±</sup>
Ascorbic acid*	20±0.55
Total phenolics**	64.37 ± 0.14
Flavonoids***	48.8 ± 2.15
DPPH inhibition %	59.42 ± 2.96
IC <sub>50</sub> (mg/ml)	36.43 ± 1.32

± Mean ± S.D. on dry weight basis

\* mg/100g

\*\* Gallic acid equivalent

\*\*\* Rutin equivalent


#### Bioactive components and antioxidant activity of Fennel seeds

Phenols and flavonoids are considered the major groups of compounds which are very important as antioxidants and having different therapeutic and protective effects on human health. Table (4) shows total phenol and total flavonoid and the bioactive components as well as the antioxidant activity of Fennel seeds.

It can be noted that Fennel seeds contained 64.37 mg/100 g total phenolics as gallic acid equivalent. On the other hand, the results shown in Table (4) indicated that flavonoids as ruitin equivalent was 48.8 mg/100 g, accordance with the results obtained in the present study the antioxidant activity (DPPH inhibition %) as well as IC<sub>50</sub> (mg/ml) of Fennel seeds were shown in Table (4) these values were 59.42 % and 36.43 mg/ml, respectively. These values are mainly due to its content of phenolic and flavonoids. These results are in agreement with those of Faudale *et al.* (2008) who found that the amounts of total phenolic in Fennel seeds ranged from 34.9 to 106.7 mg /100 g, and the total flavonoids content of the ethanol extracts from 34.1 to 95.8 mg/100 g. The antioxidant activity (DPPH inhibition %) as well as IC<sub>50</sub> (mg/ml) ranged from 46.1 to 226.3 %), (10.5 to 134.7 mg/m) respectively. Mohamad *et al.*(2011) and Shahat *et al.* (2011) were found that there is discrepancy in the values of antioxidant activity is due to existence different species of fennel. There are medicinal fennel, edible fennel and wild fennel. Different agriculture places lead to differences antioxidant activity where the antioxidant activity of the medicinal fennel was found to be higher than that of edible fennel but lower than that of wild fennel.

#### Antimicrobial activity of essential oils and plant extracts

properties of Fennel extracts and its essential oil, and their components inhibit the growth of microorganisms in food which leads to stop the deterioration of foods and foodborne pathogens on/in food its can diminution nutritional quality of the food by microorganisms which are consumption essential nutrients that are existent in



**Fig. 1** Antimicrobial activity of Fennel seeds and oil extracts using disc diffusion assay (Inhibition zone diameter in mm)

**Table 5.** microbiological properties evaluation for biscuit containing Fennel seed and its essential oil stored at room temperature for different times

product	test	time	Additive			
			0	Fennel seed		25 % Fennel seed + 2 % essential oil
				% 25	% 50	
		0	ND*	ND*	ND*	ND*
Biscuit	TVC (cfu/ml)	30 day	10x3±0.1	10x1±0.1	ND	ND
		60 day	10 <sup>2</sup> ±0.1	10x3±0.1	10x1±0.1	ND
		90 day	10 <sup>2</sup> x2±0.1	10x4±0.1	10x1±0.1	10x1±0.1

\*ND = Not Detected

the food, thereafter causes food discoloration, mustiness, biochemical changes, and toxicity compounds which can adversely affect the health of human.

Antimicrobial activity of extract of fennel seed and the essential oils are shown in Fig. 1.

The results indicated that all samples have antibacterial activity against Gram negative and Gram-positive bacteria. The effect of the sample was higher on Gram-positive bacteria *Staphylococcus aureus*, than Gram-negative bacteria *Escherichia coli*. While the oil extract was more effective than the seed extract in the diffusion assay (inhibition zone diameter in mm) it was 10, 8 and 12, 9 respectively. In addition, the higher the concentration of extract from oil or seeds the greater the diameter of the inhibition zone. When the extract concentration of oil and seeds was 20 µg / disk, the diameter of the inhibitory region was (27, 25 and 21, 20 mm) *Staphylococcus aureus* and *Escherichia coli* respectively. The same approach is applied in yeasts and mold for the concentration of the extract while the

extract has the same effect on both yeasts and mold. These results are consistent with what Anwar *et al.* (2009) got results, While these data in this study vary with results Shahat *et al.* (2011).

These results enable us to use fennel seed as natural food preserver. Fennel is recommended because it is economic value. In addition, it can be used preparation in foods and local industries as functional foods, besides its traditional uses.

**Effect of Fennel seed and its essential oil on microbial activity of stored biscuit**

Table (5) shows the microbiological properties evaluation for biscuit containing Fennel seed and its essential oil stored at room temperature for different times.

Results showed that the bacterial growth was inhibited in biscuit with increased percentage of Fennel seed and its essential oil during storage. These results are in accordance with results obtained by; Chimnoi *et al.* (2018) found that the fennel seed and its essential oil

**Table 6. Evaluation of sensory properties of biscuit during storage period at room temperature**

Time	% concentrate	color	Taste	Odor	Textures	General Acceptance
0	0	8.40 ± 0.10 <sup>a</sup>	8.32 ± 0.08 <sup>a</sup>	8.37 ± 0.15 <sup>a</sup>	8.40 ± 0.10 <sup>a</sup>	8.40 ± 0.08 <sup>a</sup>
	Fennel % 25	8.40 ± 0.10 <sup>a</sup>	8.32 ± 0.08 <sup>a</sup>	8.37 ± 0.15 <sup>a</sup>	8.40 ± 0.10 <sup>a</sup>	8.40 ± 0.08 <sup>a</sup>
	seed % 50	8.37 ± 0.15 <sup>a</sup>	8.27 ± 0.06 <sup>a</sup>	8.37 ± 0.15 <sup>a</sup>	8.37 ± 0.15 <sup>a</sup>	8.37 ± 0.15 <sup>a</sup>
	25 % Fennel seed + 2 % essential oil	8.40 ± 0.10 <sup>a</sup>	8.40 ± 0.10 <sup>a</sup>	8.40 ± 0.10 <sup>a</sup>	8.37 ± 0.15 <sup>a</sup>	8.40 ± 0.10 <sup>a</sup>
30 day	0	7.35 ± 0.20 <sup>b</sup>				
	Fennel % 25	8.40 ± 0.10 <sup>a</sup>				
	seed % 50	8.40 ± 0.10 <sup>a</sup>	8.40 ± 0.10 <sup>a</sup>	8.40 ± 0.10 <sup>a</sup>	8.37 ± 0.15 <sup>a</sup>	8.40 ± 0.15 <sup>a</sup>
	25 % Fennel seed + 2 % essential oil	8.40 ± 0.10 <sup>a</sup>	8.40 ± 0.10 <sup>a</sup>	8.40 ± 0.10 <sup>a</sup>	8.37 ± 0.15 <sup>a</sup>	8.37 ± 0.15 <sup>a</sup>
60 day	0	6.35 ± 0.20 <sup>c</sup>	6.00 ± 0.20 <sup>c</sup>			
	Fennel % 25	7.35 ± 0.20 <sup>b</sup>				
	seed % 50	8.40 ± 0.10 <sup>a</sup>				
	25 % Fennel seed + 2 % essential oil	8.40 ± 0.10 <sup>a</sup>				
90 day	0	4.00 ± 0.20 <sup>d</sup>				
	Fennel % 25	6.35 ± 0.20 <sup>c</sup>	6.00 ± 0.20 <sup>c</sup>			
	seed % 50	8.00 ± 0.10 <sup>a</sup>				
	25 % Fennel seed + 2 % essential oil	8.40 ± 0.10 <sup>a</sup>				

Numbers are mean ± standard deviation. Different superscript letters in the same column indicate mean values differ significantly ( $p < 0.05$ ).

had an effect on the membrane integrity in bacteria. Fennel seed and its essential oil work to disable the damage of the cell membrane. The level of cell damage depended on the different concentrations of the extract and the bacterial species also. Because caused the leakage of the intracellular constituents including proteins which finally results in cell death. these results make it possible to use fennel seeds as a natural preservative for food.

#### Sensory evaluation of some traditional food products

Table (6) shows sensory attributes of some traditional food products (biscuit).

It can be shown from Table (6) that there is no significant differences in the organoleptic contributes of Biscuit containing different ratios of Fennel seed and its oil comparing with the control sample at zero time. There were significant differences between the control sample and the organoleptic contributes of biscuit containing the different ratios of Fennel seed and oil during the storage periods. On the other hand, degree of

sensory properties given increased with the increase in the percentage of Fennel seed and oil with the length of storage period. Mariod, 2016 found that The oil from the Fennel seed are responsible for providing a typical flavor owing to the presence of anethole, Estragol, limonene” These constituents give a warm, aromatic and pleasing flavor to food products when used in processed foods, snacks, sauces, and various vegetables . Addition of some essential oils to foods during processing can prevent their oxidation rate and reduce oxidation effects without any changes in color and texture properties.

#### Effect of fennel and its oil in the treatment of cough and bronchitis.

Results showed that in the control group, whether the person is a smoker or not, it took more than a month to heal. The group that took doses of fennel extract and its oil, it took then 7 to 3 days to heal depending on the dose and number of times was taken during the day (Table 7).

**Table 7. Effect of fennel seed and its oil in the treatment of cough and bronchitis**

group		Dose	healing period
control	smoke	medicine	more than
		Specific dose	a 30 day
	No- smoke	medicine	30 day
		Specific dose	
Test- group	smoke	Beverage 5 g of fennel seeds	7 day
		Beverage 10 g of fennel seeds	7 day
		Beverage 15 g of fennel seeds	5 day
	No- smoke	Beverage 5 g of fennel seeds	7 day
		Beverage 10 g of fennel seeds	3 day
		Beverage 15 g of fennel seeds	3 day

Results of Choi and Hwang, 2004 study showed a significant anti-inflammatory, and analgesic activities from *F. vulgare* extract when given at dose of 200 mg/kg in mice and rats. Badgular *et al.* 2014 also reported that the fennel extract was effective in bronchial persons.

This study agreed with the study of Jahan, et al. (2015). They prepared a syrup from fennel seeds, its oil; they evaluated the syrup effect on decreasing coughing in rats. After they were exposed to a dose of SO<sub>2</sub>. Experimental rats were divided into three groups (n = 6). The first group (control), received a standard medication for cough, coding phosphate (10 mg / kg). From group 2 to 3, cough syrup was given at 0.25 ml / kg and 0.5 mg / kg body weight to the rats. After 30 minutes taking the dose, rats were exposed to sulfur dioxide for 30 seconds. The mice were then placed under observation to calculate coughing seizures for five minutes. Cough Syrup showed the decrease of cough inhibition was 63.91% and 70.64% respectively inhibition of cough frequency at 0.25mL / kg and 0.5mL / kg dose level, respectively. Highly effective anti-tussive Cough Syrup proved in the cough model caused by sulfur dioxide. Thus, Cough Syrup was useful as an alternative medicine for cough which are having a lot of side effects.

#### REFERENCES

Ahmad, B. S., T. Talou, Z. Saad, A. Hijazi, M. Cerny, H. Kanaan, A. Chokr and O. Merah. 2018. Fennel oil and by-products seed characterization and their potential Applications. *Industrial Crops & Products*. 111: 92–98.

- Al-Reza, S. M., A. Rahman, J. H. Lee and S. C. Kang. 2010. Potential roles of essential oil and organic extracts of *Zizyphus jujube* in inhibiting food-borne pathogens. *Food Chemistry*. 119: 981-986.
- Anwar, F., M. Ali, A. I. Hussain and M. Shahid. 2009. Antioxidant and antimicrobial activities of essential oil and extracts of fennel (*Foeniculum vulgare* Mill.) seeds from Pakistan. *Flavour Fragr. J.* 24: 170–176.
- Association of Official Analytical Chemists (AOAC). 2006. *Official Methods of Analytical Chemists* 20th ed. Arlington, Virginia, USA.
- Badgular, S. B., V. V. Patel and A. H. Bandivdekar. 2014. *Foeniculum vulgare* Mill: A Review of Its Botany, Phytochemistry, Pharmacology, Contemporary Application, and Toxicology, Hindawi Publishing Corporation BioMed Research International.1-33.
- Bajpai, V. K., K. H. Baek and S. C. Kang. 2012. Control of Salmonella in foods by using essential oils: a review. *Food Research International*. 45: 722-734.
- Barros, L., A. M. Carvalho and I. C. F. R. Ferreira. 2010. "The nutritional composition of fennel (*Foeniculum vulgare*): shoots, leaves, stems and inflorescences," *LWT: Food Science and Technology*. 43: 814–818.
- Bukhari, H., A. Shehzad, K. Saeed, M. S. Butt, S. Tanveer, T. Iftikhar and U. S. Din. 2014. Compositional profiling of fennel seed. *Pak. J. Food Sci.* 24: 132-136.
- Burkhardt, A., H. Y. Sintim, A. Gawde, C. L. Cantrell, T. Astatkie, V. D. Zheljzkov, V. Schlegel. 2015. Method for attaining fennel (*Foeniculum vulgare* Mill.) seed oil fractions with different composition and antioxidant capacity. *Journal of Applied Research on Medicinal and Aromatic Plants*. 2: 87–91.

- Corona, M., M. A. Cabrera, O. Santiago, E. Gonzalez, I. Palacios and J. Herrera. 2008. Activity against Drug Resistant-Tuberculosis Strains of Plants used in Mexican Traditional Medicine to treat Tuberculosis and Other Respiratory Diseases, *Phytotherapy Research*. 22: 82– 85.
- Chaturvedi, S., P.K. Sharma, V.K. Garg, M. Bansal. 2011. Role of nutraceuticals in health promotion. *Inter. J. PharmTech. Res.* 3: 442-448.
- Chimnoi, N., N. Reuk-ngam, P. Chuysinuan, P. Khlaychan, K. Nisachon, C. Daranee, W. Thamniyom, S. Klayraunge, C. Mahidol and S. Techasakul. 2018. Characterization of essential oil from *Ocimum gratissimum* leaves: Antibacterial and mode of action against selected gastroenteritis pathogens. 118: 290-300.
- Choi, E., J. Hwang. 2004. Antiinflammatory, analgesic and antioxidant activities of the fruit of *Foeniculum vulgare*. *Fitoterapia*. 75: 557– 565.
- Damayanti, A. and E. Setyawan. 2012 Essential Oil Extraction of Fennel Seed(*Foeniculum vulgare*) Using Steam, *Int. J. Sci. Eng.* 3:12-14.
- Diao, W. R., Q. Hu, H. Zhang, J. Xu. 2014. Chemical composition, antibacterial activity and mechanism of action of essential oil from seeds of fennel (*Foeniculum vulgare* Mill.). *Food Control*. 35: 109-116.
- Faudale, M., F. Viladomat, J. Bastida, F. Poli and C. Codina. 2008. Antioxidant Activity and Phenolic Composition of Wild, Edible, and Medicinal Fennel from Different Mediterranean Countries. *J. Agric. Food Chem.* 56:1912–1920.
- Gori, L., E. Gallo, V. Mascherini, A. Mugelli, A. Vannacci and F. Firenzuoli. 2012. Can estragole in fennel seed decoctions really be considered a danger for human health? A fennel safety update? *J. Evid-Based Complementary and Alternative Medi.* 1-10.
- He, W. and B. Huang. 2011. A review of chemistry and bioactivities of a medicinal spice: *Foeniculum vulgare*. *J. Med Plants. Res.* 5(16): 3595-3600.
- Jahan, Y., T. Mahmood, P. Bagga, A. Kumar, K. Singh and M. Mujahid. 2015. Future Prospects of Cough Treatment; Herbal Medicines v/s Modern Drugs. *IJPSR.* 6: 1000-09.
- Kaskatepe, B., M. E. Kiymaci, S. Suzuk, S. A. Erdem, S. Cesur and S.Yildiz. 2016. Antibacterial effects of cinnamon oil against carbapenem resistant nosocomial *Acinetobacter baumannii* and *Pseudomonas aeruginosa* isolates. *Industrial Crops and Products.* 81: 191–194.
- Keservani, R. K., R. K. Kesharwani, N. Vyas, S. Jain, R. Raghuvanshi, A.K. Sharma. 2010. Nutraceutical and Functional Food as Future Food. *Der Pharmacia Lettre.* 2 (1) :106-116.
- Kontogiorgis, C., G. Deligiannidou, D. H. Litina, D. Lazari, A. Papadopoulos. 2016. Antioxidant protection: The contribution of proper preparation of fennel (*Foeniculum vulgare* Mill.) beverage. *Industrial Crops and Products.* 79: 57–62.
- Koudela, M. and K. Petříková. 2008. Nutritional compositions and yield of sweet fennel cultivars – *Foeniculum vulgare* Mill. ssp. *vulgare* var. *azoricum* (Mill.) Thell. *Hort. Sci. (Prague)*. 35: 1–6.
- Malhotra, S. K.2012. Chapter 14, Fennel and fennel seed, *Handbook of herbs and spices.* 275-302.
- Mansour, S. A., T. M. Heikal, A. A. Refaie, A. H. Mossa. 2011. Antihepatotoxic activity of fennel (*Foeniculum vulgare* Mill.) essential oil against chlorpyrifos-induced liver injury in rats. *Global J. Environ. Sci. Techno.* 1- 10.
- Mariod, A. A. 2016. Effect of Essential Oils on Organoleptic (Smell, Taste, and Texture) Properties of Food. Chapter 13, *Essential Oils in Food Preservation, Flavor and Safety.* pp. 131-137.
- Mashat, B. H. 2013. Are Exposures TO Indoor Fungi Causing Specific Disease?. *Ass. Univ. Bull. Environ. Res.* 16:27-44.
- Mohamad, R. H., A. M. El-Bastawesy, M. G. Abdel-Monem, A. M. Noor, H. A. Al-Mehdar, S. M. Sharawy and M. M. El-Merzabani. 2011. Antioxidant and Anticarcinogenic Effects of Methanolic Extract and Volatile Oil of Fennel Seeds (*Foeniculum vulgare*), *J Med Food.* 14 : 986–1001
- Qiu, J., H. Li, H. Su, J. Dong, M. Luo, J. Wang, B. Leng, Y. Deng, J. Liu, X. Deng. 2012. Chemical composition of fennel essential oil and its impact on *Staphylococcus aureus* exotoxin production. *World J Microbiol Biotechnol.* 28:1399–1405.
- Rather, M. A., B. A. Dar, S. N. Sofi, B. A. Bhat and M. A. Qurishi. 2012. *Foeniculum vulgare*: a comprehensive review of its traditional use, phytochemistry, pharmacology, and safety. *Arabian Journal of Chemistry.* 9: S1574–S1583.
- Roby, M. H. H., M. A. Sarhan, K. A. Selim, K. I. Khalel. 2012. Antioxidant and antimicrobial activities of essential oil and extracts of fennel (*Foeniculum vulgare* L.) and chamomile (*Matricaria chamomilla* L.). *Industrial Crops and Products.* 44:437-445.
- Ruberto, G., M.T. Baratta, S.G. Deans and H.J.D. Dorman. 2000. Antioxidant and Antimicrobial Activity of *Foeniculum vulgare* and *Crithmum maritimum* Essential Oils. *Planta Medica.* 66: 687-693.
- Salgueiro, L., A. P. Martins and H. Correia. 2010. Raw materials: the importance of quality and safety. A review. *Flavour Fragrance Journal.* 25: 253-271.
- Shahat, A. A., A. Y. Ibrahim, S. F. Hendawy, E. A. Omer, F. M. Hammouda, F. H. Abdel-Rahmanand, , M.A. Saleh. 2011. Chemical Composition, Antimicrobial and Antioxidant Activities of Essential Oils from Organically Cultivated Fennel Cultivars, *Molecules.* 16: 1366-1377.
- Telci, I., I. Demirtas, and A. Sahin. 2009. Variation in plant properties and essential oil composition of sweet fennel (*Foeniculum vulgare* Mill.) fruits during stages of maturity. *Industrial Crops and Products.* 30: 126-130.
- USDA. 2010. National Nutrient Database for Standard Reference. US Department of Agriculture/Agriculture Research Service, Washington DC.

USDA. 2018. National Nutrient Database for Standard Reference Release 1 April, Full Report (All Nutrients) June. 17. 2019. 17-14 EDT.

Wichchukita, S. and M. O'Mahony. 2014. The 9-point hedonic scale and hedonic ranking in food science: some reappraisals and alternatives. Society of Chemical Industry. J Sci Food Agric.

## الملخص العربي

### استخدام حبوب زيت الشمر كمادة حافظة وغذاء وظيفي لإنتاج بعض الأغذية والمشروبات لتخفيف اعراض الكحة

جيهان إبراهيم صابر<sup>1</sup>، داليا حسن عشرة<sup>2</sup>

بذور الشمر وزيته العطري يحتويان على مواد فعالة لها تأثير في تخفيف أعراض الكحة والتهاب الحنجرة. كما كان لمستخلص الميثانول من بذور الشمر أنشطة مضادة للأكسدة ومضادات الميكروبات. كما أظهرت هذه الدراسة أيضاً أن البسكويت الذي يحتوي على بذور الشمر وزيته العطري يمكن أن يخزن لفترات أطول من البسكويت غير المحتوي على بذور الشمر وزيته العطري دون حدوث أي تغيرات في الخواص الحسية أو ظهور نموات ميكروبية خلال فترة التخزين، لذلك يمكننا استخدامه كمادة حافظة طبيعية. بالإضافة إلى ذلك تم قبول المنتج المُخزن والمُحتوي على نسب مختلفة من بذور الشمر وزيته العطري من قبل المحكمين. والحفاظ على خصائصه الحسية بشكل جيد. أظهرت النتائج أيضاً أن المشروب المعد من الشمر وزيته له تأثير فعال في علاج الكحة.

يحتوي بذور الشمر وزيته العطري على مكونات نشطة بيولوجيا ذات قيمة غذائية مرتفعة بالإضافة إلى تأثيرهما المضاد للميكروبات والفطريات والتخفيف من أعراض بعض الأمراض. لذلك كان الهدف من هذا البحث هو معرفة الخواص الفيزيائية لمستخلص بذور الشمر وزيته العطري وتأثيرهما على نمو الميكروبات والتخفيف من أعراض أمراض الجهاز التنفسي مثل الكحة والتهاب الحنجرة.

تم تحضير مشروب من الشمر وزيته العطري بنسب مختلفة لعلاج مجموعة من الأفراد المصابين بالكحة ومقارنتهم بالمجموعة الضابطة. وكذلك تم إعداد بسكويت يحتوي على نسب مختلفة من الشمر وزيته العطري.

أظهرت نتائج هذه الدراسة أن بذور الشمر يحتوي على نسبة مرتفعة من البروتين، الألياف الغذائية، الكربوهيدرات، المعادن، الفيتامينات والفلافونويدات. كذلك أظهرت النتائج ان